


“Susy-Inspired” Models



- Phenomenology fixed by boundary conditions to RGEs at high scale
- Depends on β -functions
- (only “Active” sparticles contribute)

Approximate Solutions:

$$m_{\tilde{q}}^2 = m_0^2 + 6.3(5.8)M_{\frac{1}{2}}^2$$

$$M_{\tilde{g}} = 2.6M_{\frac{1}{2}} = 3M_{C_1} = 6M_{N_1}$$


$$M_{\tilde{g}}, m_{\tilde{q}} \rightarrow M_{\frac{1}{2}}, m_0$$

$$m_0 < 0 \Rightarrow m_{\tilde{\nu}} = 350 \text{ GeV}$$

Explores possibilities *beyond* mSUGRA



S. Mrenna
TeV
Connections



“Susy-Inspired” Models



S. Mrenna


TeV

Connections

- $m_{\tilde{t}_1} = m_{\tilde{t}_2} = \sqrt{m_{\tilde{q}}^2 + m_t^2}$
- $M_A, \tan \beta$, Stop determine Higgs sector
- $M_A \gg M_Z \Rightarrow$ SM-like light Higgs
- Alternatively, ignore SM-like Higgs in “exact” mSUGRA
- But, typically, mSUGRA has $m_{\tilde{q}} > M_{\tilde{g}}$
- “Arbitrary” choice of slepton mass fixes leptonic BRs of C_1 (like-sign dileptons, etc)



Same in mSugra Framework



- $m_{H_1} = m_{H_2} = m_0$
- light Higgs boson highly correlated with other phenomenology

Specific Models

Low-Scale Parameter Sets

“Realistic”

Highly correlated


Not a simple high-scale picture

Recyclable

Can point to analysis improvements





S. Mrenna
TeV
Connections



Trilepton Results



S. Mrenna
TeV
Connections



- Sensitivity to variation of parameters
- m_0 and $M_{\frac{1}{2}}$ contours

- Sensitivity to variation of BRs
- Not from a fully consistent model



Model/Parameter Dependence



S. Mrenna
TeV
Connections

- Production and Decay Rates fixed by mixings in \tilde{N}, \tilde{C} sectors, squark/slepton masses

In general, heavy objects produced with central rapidity

- Efficiency fixed by mass splitting $M_{\tilde{C}_1} \simeq M_{\tilde{N}_2} - M_{\tilde{N}_1}$


Decay distribution described by phase space, except near kinematic boundaries

Show dependence on

- $M_2 - M_1$
- Choice of decay distribution
- $\tan \beta$



Parameter Tree



S. Mrenna

TeV

Connections

High Scale Models
(boundary conditions
for RGEs at high scale)




Minimal SuperGravity Paradigm

- (a) $m_L = m_E = m_Q = m_U = m_D = m_H = m_0$
- (b) $M_B = M_W = M_g = M_{\frac{1}{2}}$
- (c) $A_t = A_b = A_\tau = A_0$
- (d) $\tan \beta$
- (e) $\mu \Leftrightarrow m_{H_1}, m_{H_2}, \tan \beta$




Parameter Tree




S. Mrenna

TeV

Connections




Parameter Tree




S. Mrenna

TeV

Connections



Parameter Tree




S. Mrenna

TeV


Connections



Parameter Tree



S. Mrenna
TeV
Connections



Caveats, other Considerations



S. Mrenna
TeV
Connections

- Fast Detector Simulations are needed when varying many parameters
- More powerful results from combining different search channels

Requires a reliable description of the Standard Model

- Most powerful result:
 - Introduce generic particles with generic couplings and look *everywhere*
- Susy results can be recycled into (e.g.) Technicolor ones
- Other possibilities (turning on some or all RPV couplings, extended Susy sectors)

